

Some Topics on Frobenius–Lusztig Kernels, II

Eric Müller*

*Mathematisches Institut, Universität München, Theresienstrasse 39,
D-80333, Munich, Germany*

metadata, citation and similar papers at core.ac.uk

Received September 4, 1997

We discuss a method to classify all Hopf subalgebras of a certain class of pointed Hopf algebras including quantized universal enveloping algebras and Frobenius–Lusztig kernels. We find factor Hopf algebras of the dual quotients of the coordinate rings of quantum $GL(n)$. We consider a large class of quasitriangular structures for the Frobenius–Lusztig kernels and compute ribbon elements.

© 1998 Academic Press

Key Words: Frobenius–Lusztig kernels; half quantum groups; sub-Hopf algebras; quasitriangular structures; ribbon elements.

We continue the investigation of restricted specializations and Frobenius–Lusztig kernels from [10].

In Section 6, a new method is described to classify all Hopf subalgebras of a certain class of pointed Hopf algebras. This can be applied to \tilde{u} and U and gives the result that all Hopf subalgebras of \tilde{u} and of U are generated as algebras by certain subsets of the canonical algebra generators of \tilde{u} and U , respectively.

As an application, in Section 7 we describe all factor bialgebras of the Frobenius kernels of $\mathcal{O}_q(SL_n)$ and $\mathcal{O}_q(GL_n)$, the coordinate rings of quantum SL_n and quantum GL_n , respectively, and find their central group-like elements.

In Section 8 we define analogues of the “Quasi- \mathcal{R} -Matrix” from [6] for the Frobenius–Lusztig kernels and use it to find a class of quasitriangular structures which contains all quasitriangular structures from the classifica-

*This is part of the author’s doctoral dissertation which the author has written at Munich University under the supervision of Professor H.-J. Schneider. The author belongs to Graduiertenkolleg “Mathematik im Bereich ihrer Wechselwirkung mit der Physik” at Munich University. E-mail: emueller@rz.mathematik.uni-muenchen.de.

tion in [4]. We show that each quasitriangular structure admits a ribbon element.

6. HOPF SUBALGEBRAS OF FROBENIUS–LUSZTIG KERNELS

We show that all Hopf subalgebras of U and \tilde{u} (under some assumptions on the order of q) are generated as algebras by certain subsets of the usual algebra generators of these algebras. Note that the *normal* Hopf subalgebras have already been classified in the appendix of [1].

PROPOSITION 6.1. *Let J_1, J_2 be finite index sets and $J := J_1 \dot{\cup} J_2$. Let H be a pointed Hopf algebra over k and α_1 (resp. α_2) be subalgebras which are generated as algebras by elements t_j which are $(g_j, 1)$ -primitive (in H , where $g_j \in G(H)$) for $j \in J_1$ (resp. $j \in J_2$). Define $V := \{t_j \mid j \in J\}$. Assume the following conditions:*

(a) *The map $\alpha_1 \otimes \alpha_2 \otimes G(H) \rightarrow H$, $a_1 \otimes a_2 \otimes g \mapsto a_1 a_2 g$ is a k -linear isomorphism.*

(b) *All $(g, 1)$ -primitive elements (for $g \in G(H)$) are contained in the vector subspace of H spanned by $G(H)$ and V . For all $j \in J$ and $g \in G(H)$ we have $t_j g = \alpha_{g,j} g t_j$ where $\alpha_{g,j} \in k \setminus \{0\}$.*

(c) *For all $j \in J$ we have $\alpha_j := \alpha_{g_j, j} \neq 1$, and $g_i = g_j \Rightarrow \alpha_i \neq \alpha_j$ for distinct $i, j \in J$.*

(d) *For $s \in \{1, 2\}$, α_s is an $\mathbb{N}_0[J_s]$ -graded algebra, where the gradation is given by $|t_j| = j$ for $j \in J_s$ (that is, all relations between the algebra generators are generated by homogeneous ones). This induces a gradation on H given by $|a_1 a_2 g| := |a_1| + |a_2| \in \mathbb{N}_0[J]$ for homogeneous elements $a_1 \in \alpha_1$, $a_2 \in \alpha_2$. Denote the vector subspace of homogeneous elements of degree $\nu \in \mathbb{N}_0[J]$ in H by H_ν .*

Then the Hopf subalgebras H' are exactly the algebras generated by subsets $T \subset V$ and subgroups $G \subset G(H)$ such that $t_j \in T$ implies $g_j \in G$, as algebras. All finite-dimensional subbialgebras of H are Hopf subalgebras.

Proof. Obviously the mentioned algebras are Hopf subalgebras of H . We show that this list contains all of them. Let H' be a subbialgebra of H and $G = G(H')$. If H' is finite-dimensional, then the submonoid G of the group $G(H)$ is finite and a group. If H' is a Hopf subalgebra then G is a group, too. Let (H'_n) be the coradical filtration of H' . Now H' is pointed as a subbialgebra of H , whence H'_0 is spanned by group-like elements and H'_1 by G and (g, h) -primitive elements where $g, h \in G$ by the Taft–Wilson theorem [12] or [9, Theorem 5.4.1]. In particular, G contains g_j if $t_j \in H'$. Let $g, h \in G$ and let $s \in H'$ be (h, hg) -primitive. Then $h^{-1}s$ is

$(1, g)$ -primitive in H' and also in H , whence

$$h^{-1}s = \lambda_0(g-1) + \sum_{j \in J: g_j = g} \lambda_j t_j$$

where λ_0, λ_j belong to k . Since H' contains $h^{-1}s$ and g , it also contains the elements

$$g^r h^{-1} s g^{-r} = \lambda_0(g-1) + \sum_{j \in J: g_j = g} \alpha_j^{-r} \lambda_j t_j$$

for all $r \in \mathbb{N}_0$. By assumption, the coefficients α_j therein are distinct and not equal to 1, whence this system of linear equations for $\lambda_j t_j$ can be solved uniquely and $t_j \in H'$ if $\lambda_j \neq 0$. Let $T = H' \cap V$. Hence H'_1 is spanned by G and products of elements of T and G as vector space. Let \tilde{H} be the subalgebra of H' generated by H'_1 . Assume that there is a minimal positive integer s such that $H'_s \not\subset \tilde{H}$ and let $x \in A_s \setminus \tilde{H}$. Using the Taft–Wilson theorem, we can assume without loss of generality

$$\Delta(x) - x \otimes g - h \otimes x \in A_{s-1} \otimes A_{s-1}, \quad (21)$$

where $g, h \in G$. We can write x as a finite sum $x = \sum_{p=1}^M x_{p, \nu_p} \gamma_p$ such that for all p , $x_{p, \nu_p} \in \alpha_1 \alpha_2 \setminus \{0\}$ is homogeneous of degree $\nu_p \in \mathbb{N}_0[J]$ and $\gamma_p \in G(H)$. We show first that $\gamma_p \in G$ for all p . For $\nu \in \mathbb{N}_0[J]$ let

$$y_\nu := \sum_{p: \nu_p = \nu} x_{p, \nu_p} \gamma_p, \quad \text{hence } x = \sum_\nu y_\nu.$$

In this sum for y_ν we can assume $\gamma_p \neq \gamma_{p'}$ whenever $p \neq p'$. Now it follows from

$$\Delta H_{\nu'} \subset \sum_{\nu_1 + \nu_2 = \nu'} H_{\nu_1} \otimes H_{\nu_2} \quad (22)$$

for all $\nu' \in \mathbb{N}_0[J]$ that $\Delta(y_\nu)$ can be obtained from $\Delta(x)$ as the image of the projection of $H \otimes H$ to

$$\sum_{\nu_1 + \nu_2 = \nu} H_{\nu_1} \otimes H_{\nu_2}.$$

The contribution of $\Delta(x) - x \otimes g - h \otimes x$ to $H_\nu \otimes H_0$ is

$$\sum_{p: \nu_p = \nu} x_{p, \nu_p} \gamma_p \otimes \gamma_p - y_\nu \otimes g \text{ if } \nu \neq 0 \quad \text{and}$$

$$\sum_{p: \nu_p = 0} \gamma_p \otimes \gamma_p - \gamma_p \otimes g - h \otimes \gamma_p$$

otherwise and belongs to $\tilde{H} \otimes k[G]$ by (21). Thus $\gamma_p \in G$ for all p , using linear independence of distinct group-like elements. For each $\nu = \sum_{j \in J} \alpha_j j \in \mathbb{N}_0[J]$, call α_j the j -coordinate of ν . The elements of \tilde{H} are

sums of homogeneous elements of H ; if the homogeneous summand in H_ν does not vanish then $t_j \notin T$ implies that the j -coordinate of ν vanishes. Since the group-like elements γ_p belong to \tilde{H} , there must be a homogeneous summand $y_\nu \neq 0$ with positive j -coordinate of ν and $t_j \notin T$. Now assume that y_ν is not (g, h) -primitive. Then $\Delta(y_\nu) - y_\nu \otimes g - h \otimes y_\nu$ contains non-zero homogeneous summands in

$$\sum_{\substack{\nu_1 + \nu_2 = \nu \\ \nu_1 \neq 0, \nu_2 \neq 0}} H_{\nu_1} \otimes H_{\nu_2},$$

where either ν_1 or ν_2 has positive j -component. Equation (22) implies that these summands cannot be killed by contributions of other homogeneous summands. But then (21) implies $H'_{s-1} \not\subset \tilde{H}$ in contrary to the minimality of s . Hence x is the sum of an element $x' \in \tilde{H}$ and homogeneous (g, h) -primitive summands, the sum of which being x'' . Now $x'' = x - x'$ belongs to H' , is (g, h) -primitive, and therefore belongs to H'_1 , contrary to the definition of \tilde{H} . Hence $H' = \tilde{H}$. ■

Remark 6.2. (a) There is an analogous proof of this proposition if H is as vector space isomorphic to the tensor product of more than two subalgebras and $G(H)$.

(b) If assumption (c) of the proposition is not satisfied then the above classification contains all Hopf subalgebras H' such that H'_1 (the second term of the coradical filtration containing all group-like and skew-primitive elements) is spanned by $G(H')$ and a subset $T \subset V$ as a vector space.

Now the Hopf subalgebras of the Frobenius–Lusztig kernels can be classified:

THEOREM 6.3. Assume that q fits to $\tilde{\mathbf{f}}$ and $q^{2^{j \cdot j}} \neq 1$ for all $j \in I$. Let \tilde{G} be a group consisting of central group-like elements of $\tilde{\mathbf{u}}$ such that $\tilde{K}_i \tilde{K}_j^{-1} \notin \tilde{G}$ for all $i, j \in I$ whenever $i \neq j$. Let $\tilde{\mathbf{u}} := \tilde{\mathbf{u}} / (\{g - 1 \mid g \in \tilde{G}\})$. Elements of $\tilde{\mathbf{u}}$ are denoted as in $\tilde{\mathbf{u}}$. Then the Hopf subalgebras of $\tilde{\mathbf{u}}$ are exactly those generated as algebras by a subset $T \subset \{E_j, F_j \mid j \in I\}$ and a subgroup $G \subset G(\tilde{\mathbf{u}})$ such that

$$\forall j \in I: (E_j \in T \vee F_j \in T \Rightarrow \tilde{K}_j \in G).$$

If $\tilde{\mathbf{u}}$ is finite-dimensional then all subbialgebras are Hopf subalgebras and of this type.

Proof. We establish the assumptions of Proposition 6.1. $\tilde{\mathbf{u}}$ is spanned by group-likes and $(1, \tilde{K}_i)$ -primitive elements $E_i, \tilde{K}_i F_i$ for all $i \in I$ as algebra. The statement for the skew-primitive elements follows from [10, 3.1]; moreover the commutation relations with group-like elements are satisfied

which implies (b). Together with the triangular decomposition, this gives (a), and the gradations on \tilde{u}^+ and \tilde{u}^- yield (d). Moreover for all $j \in I$

$$\tilde{K}_j E_j \tilde{K}_j^{-1} = q^{j \cdot j} E_j, \quad \tilde{K}_j (\tilde{K}_j F_j) \tilde{K}_j^{-1} = q^{-j \cdot j} \tilde{K}_j F_j.$$

The assumption on \tilde{G} and $q^{2j \cdot j} \neq 1$ for all $j \in I$ imply assumption (c). ■

THEOREM 6.4. *Let (I, \cdot) be an arbitrary Cartan datum. If the elements \tilde{K}_i in \mathbf{U} for $i \in I$ are pairwise distinct, e.g., if the root datum is Y -regular [7, 2.2.2], then the Hopf subalgebras U of \mathbf{U} are just those which are generated as algebras by a subset $T \subset \{E_j, F_j \mid j \in I\}$ and a subgroup $G \subset G(\mathbf{U})$ such that*

$$\forall j \in I: (E_j \in T \vee F_j \in T \Rightarrow \tilde{K}_j \in G).$$

Proof. We establish the assumptions of Proposition 6.1. \mathbf{U} is generated by group-likes and $(1, \tilde{K}_i)$ -primitive elements $E_i, \tilde{K}_i F_i$ for all $i \in I$ as algebra. The statement for the skew-primitive elements follows from [13]; moreover the commutation relations with group-like elements are satisfied which implies (b). Together with the triangular decomposition, this gives (a), and the gradations on \mathbf{U}^+ and \mathbf{U}^- yield (d). Moreover for all $j \in I$

$$\tilde{K}_j E_j \tilde{K}_j^{-1} = v^{j \cdot j} E_j, \quad \tilde{K}_j (\tilde{K}_j F_j) \tilde{K}_j^{-1} = v^{-j \cdot j} \tilde{K}_j F_j.$$

Now assumption (c) follows because v is not a root of unity. ■

Remark 6.5. If the determinant of the Cartan matrix is coprime to the order of q , then it is easy to show that $\tilde{K}_i \tilde{K}_j^{-1}$ is not central for $i \neq j$ (cf. [10, 4.3 (c)]). Hence in this case there are no restrictions for \tilde{G} in Theorem 6.3.

7. QUOTIENTS OF THE DUAL FROBENIUS–LUSZTIG KERNELS OF QUANTUM $GL(n)$ AND $SL(n)$

We recall some definitions from Sections 2 and 3 of [15], but we use opposite comultiplication for the quantized enveloping algebra in accordance with [6] and the other parts of this paper and opposite multiplication on the quantized matrix algebra which just means to replace q by its inverse. Let k be a field and $q \in k \setminus \{0\}$.

DEFINITION 7.1. Fix an integer $n > 1$, define a Cartan datum (I, \cdot) by $I := \{1, \dots, n-1\}$ and

$$i \cdot j = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } |i - j| = 1, \\ 0 & \text{otherwise,} \end{cases}$$

and a root datum of type (I, \cdot) by $X = Y = \mathbb{Z}^n$ and $\langle (a_j), (b_j) \rangle = \sum_{j=1}^n a_j b_j$ for $a_j, b_j \in \mathbb{Z}$, and the imbeddings $I \rightarrow X$, $I \rightarrow Y$ are given by

$$j \mapsto (0, \dots, 0, \overset{j}{\underbrace{1, -1}}, \dots, 0).$$

The quantized enveloping algebra attached to this root datum will be called $U(GL_q(n))$; the Hopf subalgebra generated by $E_i, F_i, K_i^{\pm 1}$ for $i \in I$ will be called $U(SL_q(n))$. The *quantized matrix algebra* $A(M_q(n))$ is generated by the elements α_{ij} as algebra for $1 \leq i, j \leq n$, subject to the relations

$$\begin{aligned} \alpha_{is} \alpha_{js} &= q \alpha_{js} \alpha_{is}, & \alpha_{si} \alpha_{sj} &= q \alpha_{sj} \alpha_{si} \text{ if } i < j, \\ \alpha_{it} \alpha_{js} &= \alpha_{js} \alpha_{it}, & \alpha_{is} \alpha_{jt} - \alpha_{jt} \alpha_{is} &= (q - q^{-1}) \alpha_{it} \alpha_{js} \text{ if } i < j, s < t. \end{aligned}$$

A bialgebra structure is given by

$$\Delta(\alpha_{ij}) = \sum_{s=1}^n \alpha_{is} \otimes \alpha_{sj}, \quad \varepsilon(\alpha_{ij}) = \delta_{ij}$$

for $1 \leq i, j \leq n$. Let $s \leq n$, $\mathbf{I} := \{i_1, i_2, \dots, i_s\}$, and $\mathbf{J} := \{j_1, j_2, \dots, j_s\}$ where $1 \leq i_1 < i_2 < \dots < i_s \leq n$ and $1 \leq j_1 < j_2 < \dots < j_s \leq n$. Define the *quantum subdeterminants*

$$\det_{\mathbf{J}}^{\mathbf{I}} := \sum_{\sigma \in \text{Sym}(s)} (-q)^{l(\sigma)} \alpha_{i_1 j_{\sigma(1)}} \alpha_{i_2 j_{\sigma(2)}} \cdots \alpha_{i_s j_{\sigma(s)}}$$

(for each permutation $\sigma \in \text{Sym}(s)$ let $l(\sigma)$ denote the length of σ). In particular define $X_{ij} := \det_{\mathbf{I}}^{\mathbf{I}}$ for $\mathbf{I} = \{i, i+1, \dots, j\}$ and $\det_q := X_{1n}$, the *quantum determinant*. The localization $A(GL_q(n)) := A(M_q(n))[\det_q^{-1}]$ and the quotient $A(SL_q(n)) := A(GL_q(n))/(\det_q - 1)$ are Hopf algebras. Their elements will be denoted like the elements of $A(M_q(n))$. There is a Hopf pairing $\langle \cdot, \cdot \rangle: U(GL_q(n)) \times A(GL_q(n)) \rightarrow k$ which induces a Hopf pairing of $U(SL_q(n))$ and $A(SL_q(n))$ and is given on the algebra generators by

$$\langle K_{(a_i)}, \alpha_{ij} \rangle = \delta_{ij} q^{a_i}, \quad \langle E_s, \alpha_{ij} \rangle = \delta_{si} \delta_{s+1, j}, \quad \langle F_s, \alpha_{ij} \rangle = \delta_{s+1, i} \delta_{sj}, \quad (23)$$

for $(a_i) \in Y$, $s \in I$, $1 \leq i, j \leq n$.

We can now define the dual Frobenius–Lusztig kernels.

DEFINITION 7.2. Let q be a root of unity of order N and let ℓ be the order of q^2 . Define

$$\begin{aligned}\tilde{a} &:= A(GL_q(n)) / \left(\left\{ \alpha_{ss}^N - 1, \alpha_{ij}^{\ell} \mid i \neq j \right\} \right), \\ \tilde{a}' &:= A(SL_q(n)) / \left(\left\{ \alpha_{ss}^N - 1, \alpha_{ij}^{\ell} \mid i \neq j \right\} \right),\end{aligned}$$

where $s, i, j \in \{1, \dots, n\}$. Let \tilde{u} and \tilde{u}' be the Frobenius–Lusztig kernels of $U(GL_q(n))$ and $U(SL_q(n))$. Define

$$\begin{aligned}\tilde{u} &:= \tilde{u} / \left(\left\{ K_{\mu}^N - 1 \mid \mu \in Y \right\} \right), \\ \tilde{u}' &:= \tilde{u}' / \left(\left\{ K_i^N - 1 \mid i \in I \right\} \right).\end{aligned}$$

Note that \tilde{u}, \tilde{u}' correspond to $U^{[\ell]}(GL_q(n))$ and $U^{[\ell]}(SL_q(n))$ from [15, Chap. 12], respectively.

Then we have by [14, Proposition 6.3]:

THEOREM 7.3. *The Hopf pairing from (23) induces isomorphisms $\tilde{u} \cong \tilde{a}^*$ and $\tilde{u}' \cong (\tilde{a}')^*$.*

Assume that q is a root of unity and $q^4 \neq 1$. Now we interpret the duals of the Hopf subalgebras of \tilde{u} and \tilde{u}' as factor Hopf algebras of \tilde{a} and \tilde{a}' .

THEOREM 7.4. *Let $\mathcal{T} := \{E_i, F_i \mid i \in I\}$. For each subset $T \subset \mathcal{T}$ define*

$$\begin{aligned}W_T &= \{i \in I \mid E_i \in T \vee F_i \in T\}, \\ P_T &= \{\alpha_{ij} \mid \exists h \in I: (E_h \notin T \wedge i \leq h < j) \vee (F_h \notin T \wedge j \leq h < i)\}, \\ G_T &= \{X_{ij} \mid 1 \leq i \leq j \leq n, i-1 \notin W_T \wedge j \notin W_T\}.\end{aligned}$$

Then each bi-ideal \mathcal{J} of \tilde{a} is a Hopf ideal and there is a subset $T \subset \mathcal{T}$ and a set G' of monomials in G_T such that \mathcal{J} is generated by P_T and $\{g - 1 \mid g \in G'\}$ as a two-sided ideal.

Proof. Clearly the assumptions of Theorem 6.3 are satisfied. All subalgebras of the finite-dimensional Hopf algebra \tilde{u} are Hopf algebras, whence by Theorem 7.3 all bi-ideals are Hopf ideals. For each bi-ideal \mathcal{J} of \tilde{a} define $T = \mathcal{T} \cap (\tilde{a}/\mathcal{J})^*$. For fixed T denote the two-sided ideal of \tilde{a} generated by P_T as \mathcal{J}_T . Direct calculation shows that \mathcal{J}_T is a bi-ideal (and Hopf ideal). The rest of the proof consists of several steps.

(a) The canonical projection $\tilde{a} \rightarrow \tilde{a}/\mathcal{J}_T$ maps the elements of G_T to central group-like elements.

Proof. Fix $r, s \in \{0, \dots, n\} \setminus W_T$, $r \leq s$. Define $M_1 = \{1, \dots, r\}$, $M_2 = \{r+1, \dots, s\}$, $M_3 = \{s+1, \dots, n\}$. Consider the “blocks” consisting of the elements α_{ij} , where i, j are both in M_1 or in M_2 or in M_3 . All other algebra generators vanish in \tilde{a}/\mathcal{J}_T , whence elements of different blocks commute in \tilde{a}/\mathcal{J}_T and the subalgebras generated by all elements of a block are isomorphic to quotients of quantized matrix algebras, whence $\bar{X}_{1,r}$, $\bar{X}_{r+1,s}$, and $\bar{X}_{s+1,n}$ are central group-like elements of \tilde{a}/\mathcal{J}_T .

(b) All ideals mentioned above are Hopf ideals, because by part (a), the corresponding factor algebras are quotients of the Hopf algebra \tilde{a}/\mathcal{J}_T by central group-like elements.

(c) The Hopf subalgebra $(\tilde{a}/(P_T \cup \{g-1 \mid g \in G'\}))^*$ of \tilde{u} contains T and all elements K_h where $h \in W_T$.

Proof. Using the duality in Theorem 7.3, it suffices to prove that for any element x of these and any element y of the ideal we have $\langle x, y \rangle = 0$. Since x is skew-primitive or group-like, it suffices to prove this on the generators of the ideal. We only discuss the most difficult computation: $\langle x, X_{r+1,s} - 1 \rangle = 0$ for $r, s \in \{0, \dots, n\} \setminus W_T$. We have

$$\begin{aligned} \langle K_h, X_{r+1,s} - 1 \rangle &= \langle K_h, \alpha_{r+1,r+1} \rangle \cdots \langle K_h, \alpha_{ss} \rangle - 1 \\ &= q^{\delta_{h+1,r+1} - \delta_{h,s}} - 1 = 0. \end{aligned}$$

Moreover $\langle E_h, X_{r+1,s} - 1 \rangle = \langle E_h, X_{r+1,s} \rangle = 0$ because $X_{r+1,s}$ consists of the summand $\alpha_{r+1,r+1} \cdots \alpha_{ss}$ each factor of which being in the kernel of $\langle E_h, \cdot \rangle$ and summands with at least two elements which are not in the diagonal and hence are in the kernels of $\langle K_h, \cdot \rangle$ and $\langle 1, \cdot \rangle$. A similar proof shows $\langle F_h, X_{r+1,s} \rangle = 0$.

(d) We have $\mathcal{J} \cap (\tilde{a}/\mathcal{J}_T)^* = T$, because the inclusion \supset has been shown in part (c), and \subset is trivial (because $\langle E_i, \alpha_{i,i+1} \rangle = 1 = \langle F_i, \alpha_{i+1,i} \rangle$).

(e) We have $G(\tilde{u}) \subset (\tilde{a}/\mathcal{J}_T)^*$, since the maps $\langle K_\mu, \cdot \rangle$ for $\mu \in Y$ are characters and $\langle K_\mu, \alpha_{ij} \rangle = 0$ for $\alpha_{ij} \in P_T$.

(f) For each $g \in G_T$, the order of $g + \mathcal{J}_T$ in \tilde{a}/\mathcal{J}_T is N .

Proof. By (e), $(\tilde{a}/\mathcal{J}_T)^*$ contains all group-like elements of \tilde{u} , and \tilde{a}/\mathcal{J}_T is dual to the subalgebra of \tilde{u} generated by T and $G(\tilde{u})$, because this is the biggest Hopf subalgebra H of \tilde{u} such that $\mathcal{J} \cap H = T$ (by Theorem 6.3). Fix $r, s \in \{0, \dots, n\} \setminus W_T$, where $r \leq s$. Since \tilde{a}/\mathcal{J}_T is dual to the Hopf subalgebra H of \tilde{u} generated by T and $G(\tilde{u})$ as algebra, we have to

compute the smallest positive integer z such that $\langle x, X_{r+1,s}^z - 1 \rangle = 0$ or $\langle x, X_{r+1,s}^z \rangle = \varepsilon(x)$ for all $x \in H$. Since $\bar{X}_{r+1,s}^z$ is group-like, the linear form $\langle \cdot, X_{r+1,s}^z \rangle$ is a character on H , and it suffices to prove the assertion for algebra generators of H . We have $\langle E_i, X_{r+1,s}^z \rangle = \varepsilon(E_i)$, $\langle F_i, X_{r+1,s}^z \rangle = \varepsilon(F_i)$ for all $i \in I$ by part (c). For all $(a_j) \in Y$ and $t \in \mathbb{N}$ we have

$$\langle K_{(a_j)}, X_{r+1,s}^t \rangle = \left(\langle K_{(a_j)}, \alpha_{r+1,r+1} \rangle \cdots \langle K_{(a_j)}, \alpha_{ss} \rangle \right)^t = q^{t(a_{r+1} + \cdots + a_s)}.$$

Hence the order is N .

(g) Let \mathcal{J} be a Hopf ideal of \tilde{a} and let $T := \mathcal{J} \cap (\tilde{a}/\mathcal{J})^*$. It follows from Theorem 6.3 that $(\tilde{a}/\mathcal{J})^* \subset \tilde{u}$ is generated by T and group-like elements as algebra. Assume

$$\{0, \dots, n\} \setminus W_T = \{r(1), r(2), \dots, r(s+1)\}$$

where $0 = r(1) < r(2) < \cdots < r(s+1) = n$. Define $e_j = (0, \dots, 1, \dots, 0) \in Y$ (1 at the j th coordinate) for $1 \leq j \leq n$. Since $(\tilde{a}/\mathcal{J})^*$ contains all elements $K_i = K_{e_i} K_{e_{i+1}}^{-1}$ where $i \in W_T$, these subgroups of $G((\tilde{a}/\mathcal{J})^*)$ correspond exactly to the subgroups of the group R_u generated by $K_{e_{r(1)+1}}, K_{e_{r(2)+1}}, \dots, K_{e_{r(s)+1}}$. By definition of \tilde{u} , R_u is isomorphic to $(\mathbb{Z}/N\mathbb{Z})^s$. Since there is a one-to-one correspondence between Hopf subalgebras of \tilde{u} and factor Hopf algebras of \tilde{a} , it suffices to prove that there are no more than the Hopf algebras described in (c). By part (f), the central group-likes $X_{r(t)+1, r(t+1)} + \mathcal{J}_T$, where $1 \leq t \leq s$ in \tilde{a}/\mathcal{J}_T , have order N ; they generate a group called R_a . The pairing between \tilde{u} and \tilde{a} induces a pairing between R_u and R_a given by the bi-character

$$\langle K_{e_{r(i)+1}}, X_{r(j)+1, r(j+1)} + \mathcal{J}_T \rangle = q^{\delta_{ij}}.$$

Since q has order N , there are no more relations between the commuting generators $X_{r(t)+1, r(t+1)} + \mathcal{J}_T$, and R_a is isomorphic to $(\mathbb{Z}/N\mathbb{Z})^s$. This duality gives a one-to-one correspondence between the subgroups of the finite group R_u and the quotients of R_a . Hence we have found all factor Hopf algebras of \tilde{a} .

COROLLARY 7.5. *Since \tilde{a}' is a factor Hopf algebra of \tilde{a} , the factor Hopf algebras of \tilde{a}' are exactly those, whose quantum determinant is 1.*

We determine the central group-like elements of these factor Hopf algebras.

THEOREM 7.6. *Let \mathcal{J} be a Hopf ideal of \tilde{a} and $T := \mathcal{J} \cap (\tilde{a}/\mathcal{J})^*$. The central group-like elements of \tilde{a}/\mathcal{J} are exactly the elements*

$$X_{r(1)+1, r(2)}^{\beta_1} \cdots X_{r(s)+1, r(s+1)}^{\beta_s} + \mathcal{J},$$

where $\{0, \dots, n\} \setminus W_T = \{r(1), r(2), \dots, r(s+1)\}$, where $0 = r(1) < r(2) < \cdots < r(s+1) = n$, and $\beta_1, \dots, \beta_s \in \mathbb{N}$.

Proof. Obviously the mentioned elements are central and group-like. Their number is the order of R'_a , a quotient of R_a from Theorem 7.4(g). Let R'_u be the corresponding subgroup of R_u from Theorem 7.4(g). Central group-like elements of \tilde{a}/\mathcal{I} correspond to algebra homomorphisms $\phi: \tilde{u} \rightarrow k$ such that for all $\psi \in \text{Hom}(\tilde{u}, k)$: $\psi * \phi = \phi * \psi$. This gives $\phi(x) = 0$ for $x \in T$ and $\phi(\tilde{K}_i) = 1$ if $E_i \in T$ or $F_i \in T$. Therefore these homomorphisms correspond to characters of R'_u . Their number is the order of R'_u , because R'_u is finite and abelian. Finally it follows from $|R'_u| = |R'_a|$ that there are no central group-like elements in addition to the elements mentioned above. ■

8. QUASITRIANGULAR STRUCTURES

Let q be a root of unity. The aim of this section is to construct a Quasi- \mathcal{R} -matrix for the Frobenius–Lusztig kernels imitating the procedure in [6, Chap. 4]. The Quasi- \mathcal{R} -matrix gives rise to quasitriangular structures. In particular, all quasitriangular structures from the classification in [4] will be described explicitly. Let \tilde{u}^\diamond be the Frobenius–Lusztig kernel obtained for q^{-1} instead of q . Then \tilde{u} and \tilde{u}^\diamond can be identified as sets.

DEFINITION 8.1. There is a unique isomorphism of algebras

$$^-: \tilde{u} \rightarrow \tilde{u}^\diamond, E_i \mapsto E_i, \quad F_i \mapsto F_i, \quad K_\mu \mapsto K_{-\mu}$$

for all $i \in I$, $\mu \in Y$ [6, 3.1.12]; the map $^- \otimes -: \tilde{u} \otimes \tilde{u} \rightarrow \tilde{u}^\diamond \otimes \tilde{u}^\diamond$ is a well-defined linear algebra isomorphism. Then we can define in analogy to [7, 4.1.1], $\bar{\Delta}: \tilde{u} \rightarrow \tilde{u} \otimes \tilde{u}$, $x \mapsto (- \otimes -) \Delta(\bar{x})$.

8.1. The Quasi- \mathcal{R} -Matrix of \tilde{u}

Assume that (I, \cdot) is of finite type, that q fits to $\tilde{\mathbf{f}}$, and $q_i^2 \neq 1$ for all $i \in I$. We take the original definition $B_i = (1 - q_i^{-2})^{-1}$. The proof of the following theorem can be repeated word by word (after replacing v by q) from [6, Theorem 4.1.2], because it only uses that (\cdot, \cdot) is non-degenerate (cf. [10, 2.16]).

THEOREM 8.2. (a) *There is a unique family of elements $\Theta_\nu \in \tilde{u}_\nu^- \otimes \tilde{u}_\nu^+$ (with $\nu \in \mathbb{N}_0[I]$) such that $\Theta_0 = 1 \otimes 1$ and $\Theta = \sum_\nu \Theta_\nu$ satisfies $\Delta(x)\Theta = \Theta\bar{\Delta}(x)$ for all $x \in \tilde{u}$ (identity in $\tilde{u} \otimes \tilde{u}$).*

(b) *Let B be a k -basis of $\tilde{\mathbf{f}}$ such that $B_\nu = B \cap \tilde{\mathbf{f}}_\nu$ is a basis of $\tilde{\mathbf{f}}_\nu$ for any ν . Let $\{b^* \mid b \in B_\nu\}$ be the basis of $\tilde{\mathbf{f}}_\nu$ dual to B_ν under (\cdot, \cdot) . We have*

$$\Theta_\nu = (-1)^{\text{tr } \nu} q_\nu \sum_{b \in B_\nu} b^- \otimes b^{*+} \in \tilde{u}_\nu^- \otimes \tilde{u}_\nu^+,$$

where $q_\nu = \prod_i q_i^{\nu_i}$ for $\nu = \sum_i \nu_i i \in \mathbb{Z}[I]$.

Remark 8.3. (a) The element Θ is called Quasi- \mathcal{R} -matrix (cf. [6, Paragraph 4.1.4]).

(b) Since Θ is unique, it is independent of the actual choice of basis.

(c) Since $\tilde{\mathbf{f}}$ is finite-dimensional, it is not necessary to consider a completion of $\tilde{\mathbf{u}} \otimes \tilde{\mathbf{u}}$ with respect to a certain topology (cf. [6, Paragraph 4.1.1]).

Here are some properties of the Quasi- \mathcal{R} -matrix. For an element $P = \sum_i x_i \otimes y_i \in V \otimes V$, where V is a vector space, we shall denote the elements $\sum_i x_i \otimes y_i \otimes 1, \sum_i 1 \otimes x_i \otimes y_i, \sum_i x_i \otimes 1 \otimes y_i$ by P^{12}, P^{23}, P^{13} , respectively [6, Paragraph 4.2.1].

PROPOSITION 8.4. *We have $\Theta\bar{\Theta} = \bar{\Theta}\Theta = 1 \otimes 1$ in $\tilde{\mathbf{u}} \otimes \tilde{\mathbf{u}}$.*

Proof. Compare [6, Corollary 4.1.3]. ■

PROPOSITION 8.5. *For all $\nu \in \mathbb{N}_0[I]$ we have*

$$\begin{aligned} (\Delta \otimes \text{id})\bar{\Theta}_\nu &= \sum_{\nu' + \nu'' = \nu} \bar{\Theta}_{\nu'}^{13} (1 \otimes \tilde{K}_{-\nu'} \otimes 1) \bar{\Theta}_{\nu''}^{23}, \\ (\text{id} \otimes \Delta)\bar{\Theta}_\nu &= \sum_{\nu' + \nu'' = \nu} \bar{\Theta}_{\nu'}^{13} (1 \otimes \tilde{K}_{\nu'} \otimes 1) \bar{\Theta}_{\nu''}^{12}. \end{aligned}$$

Proof. This is similar to [6, Proposition 4.2.3]. ■

Remark 8.6. The lemma in [10, 2.9] contains a more detailed description. Reference [10, Eq. (7)] gives

$$\Theta = \sum_{\mathbf{c}} (-1)^{\text{tr}|b_{\mathbf{c}}|} q_{|b_{\mathbf{c}}|} b_{\mathbf{c}}^- \otimes \tilde{b}_{\mathbf{c}}^+.$$

The Quasi- \mathcal{R} -matrix is invariant under Hopf algebra automorphisms:

PROPOSITION 8.7. *Assume that there are no weak diagram antimorphisms of (I, \cdot) (e.g., if it is irreducible). Let $\phi: \tilde{\mathbf{u}} \rightarrow \tilde{\mathbf{u}}$ be an automorphism. Then $(\phi \otimes \phi)(\Theta) = \Theta, (\phi \otimes \phi)\bar{\Theta} = \bar{\Theta}$.*

Proof. It follows from [10, 5.14] and the only exceptional case in [10, 5.15] that irreducible Cartan data of finite type do not admit weak diagram antimorphisms. By [10, 5.14] there are coefficients $a_i \in k \setminus \{0\}$ for $i \in I$ and a weak diagram automorphism σ of I such that

$$\phi(E_i) = a_i E_{\sigma(i)}, \quad \phi(F_i) = q_i q_{\sigma(i)}^{-1} a_i^{-1} F_{\sigma(i)}, \quad \phi(\tilde{K}_i) = \tilde{K}_{\sigma(i)}.$$

In particular, $(\phi \otimes \phi)(\Theta) \in \tilde{\mathbf{u}}^- \otimes \tilde{\mathbf{u}}^+$. If we can prove that

$$\bar{\Delta}\phi(x) = (\phi \otimes \phi)\bar{\Delta}(x) \tag{24}$$

for all $x \in \tilde{\mathbf{u}}$, we have $\Delta(\phi(x))(\phi \otimes \phi)(\Theta) = (\phi \otimes \phi)(\Theta)\bar{\Delta}\phi(x)$, and the uniqueness of Θ in Theorem 8.2 implies the assertion for Θ and $\bar{\Theta}$,

because $\bar{\Theta}$ is the inverse of Θ . Now we prove (24). Since the maps on both sides are k -algebra homomorphisms, it suffices to check (24) on the algebra generators. We have

$$\bar{\Delta}\phi(E_i) = a_i(E_{\sigma(i)} \otimes 1 + \tilde{K}_{\sigma(i)}^{-1} \otimes E_{\sigma(i)}) = (\phi \otimes \phi)\bar{\Delta}(E_i),$$

and a similar computation holds for F_i . For each $\mu \in Y$ we have $\bar{\Delta}(K_\mu) = K_\mu \otimes K_\mu$, whence (24) holds for $x = K_\mu$. ■

Remark 8.8. The proof of Proposition 8.7 can be applied to the situation of the Quasi- \mathcal{R} -matrix of \mathbf{U} , because the automorphisms of \mathbf{U} are known (cf. [2, Theorem C]).

8.2. Quasitriangular Structures

We recall the definition of a quasitriangular structure.

DEFINITION 8.9. Let H be a Hopf algebra. An invertible element R of $H \otimes H$ is called *quasitriangular structure*, if

$$\begin{aligned} \Delta^{\text{op}}(x)R &= R\Delta(x) \text{ for all } x \in H, & (\Delta \otimes \text{id})R &= R^{13}R^{23}, \\ (\text{id} \otimes \Delta)R &= R^{13}R^{12}. \end{aligned} \quad (25)$$

Notation 8.10. (a) For $\nu = \sum_i \nu_i i \in \mathbb{Z}[I]$ let $[\nu] = \sum_i \nu_i \frac{i \cdot i}{2}$.

(b) In the Frobenius–Lusztig kernels there are some extra relations between the group-like elements, namely $\tilde{K}_i^{2l_i} = 1$ for $i \in I$, where l_i is the order of q_i^2 . These are all relations (cf. [7, Theorem 8.3(ii)]). Let $Y' \subset Y$ be the subgroup generated by $2l_i[i]$ for $i \in I$. Define $\tilde{Y} = Y/Y'$.

THEOREM 8.11. Let $G \subset \{\mu \in \tilde{Y} \mid K_\mu \text{ central}\}$ be a subgroup. Let Y_1, Y_2 be finite subgroups of \tilde{Y}/G which contain the images of $[\nu] \in Y$. In the following $\mu, \tilde{\mu}, \mu_1, \mu_2$ and $\rho, \tilde{\rho}, \rho_1, \rho_2$ denote elements of Y_1 and Y_2 , respectively. The element $R = R_0 \bar{\Theta}$ where $R_0 = \sum_{\mu, \rho} f(\mu, \rho) K_\mu \otimes K_\rho$ is a quasitriangular structure of $\tilde{u} = \tilde{u}/(\{K_\mu - 1 \mid \mu \in G\})$ if and only if for all $i \in I$ and $\mu, \rho, \tilde{\mu}, \tilde{\rho}$,

$$f(\mu + [i], \rho) = f(\mu, \rho)q^{-\langle \rho, i' \rangle}, \quad f(\mu, \rho + [i]) = f(\mu, \rho)q^{-\langle \mu, i' \rangle} \quad (26)$$

$$\begin{aligned} \sum_{\substack{\rho_1, \rho_2 \in Y_2 \\ \rho_1 + \rho_2 = \rho}} f(\mu, \rho_1)f(\tilde{\mu}, \rho_2) &= \delta_{\mu, \tilde{\mu}}f(\mu, \rho), \\ \sum_{\substack{\mu_1, \mu_2 \in Y_1 \\ \mu_1 + \mu_2 = \mu}} f(\mu_1, \rho)f(\mu_2, \tilde{\rho}) &= \delta_{\rho, \tilde{\rho}}f(\mu, \rho), \end{aligned} \quad (27)$$

$$\sum_{\mu} f(\mu, \rho) = \delta_{\rho, 0} \vee \sum_{\rho} f(\mu, \rho) = \delta_{\mu, 0}. \quad (28)$$

Condition (28) follows from (26) and (27) if there exists $l \in k \setminus \{0\}$ such that $f(\mu, 0) = f(0, \rho) = l$ for all μ, ρ . There are conditions on the order of q : For all μ, ρ for which there exist $\tilde{\mu}, \tilde{\rho}$ such that $f(\mu, \tilde{\rho}) \neq 0, f(\tilde{\mu}, \rho) \neq 0$ we have

$$q^{2l_i \langle \mu, i' \rangle} = q^{2l_i \langle \rho, i' \rangle} = 1. \quad (29)$$

If this condition is satisfied then f is well-defined on the preimages of $Y_1 \times Y_2$ under $\tilde{Y} \rightarrow \tilde{Y}/G$.

Proof. It follows from (26) that $f(\mu + 2l_i[i], \rho) = q^{-2l_i \langle \rho, i' \rangle} f(\mu, \rho)$. These are the only relations in \tilde{u} between the commuting group-like elements. We check the conditions in Definition 8.9.

(a) $\Delta^{\text{op}}(x)R = R\Delta(x)$ or $\Delta^{\text{op}}(x)R_0 = R_0\bar{\Delta}(x)$ for all $x \in \tilde{u}$. The set of elements of \tilde{u} which satisfy this condition is a subalgebra of \tilde{u} containing all K_μ . Hence, in order for this set to be equal to \tilde{u} , it is necessary and sufficient that it contains E_i, F_i for all $i \in I$. We check the condition for $x = E_i$.

$$(1 \otimes E_i + E_i \otimes \tilde{K}_i)R_0 = R_0(E_i \otimes 1 + \tilde{K}_{-i} \otimes E_i).$$

With respect to the gradation $\tilde{u}_{(\nu, \gamma)} = t(\tilde{\mathbf{f}}_\nu \otimes \tilde{u}^0 \otimes \tilde{\mathbf{f}}_\gamma)$, the elements $(1 \otimes E_i)R_0$ and $R_0(\tilde{K}_{-i} \otimes E_i)$ belong to $\tilde{u}_{(0,0)} \otimes \tilde{u}_{(i,0)}$ and $(E_i \otimes \tilde{K}_i)R_0, R_0(E_i \otimes 1)$ to $\tilde{u}_{(i,0)} \otimes \tilde{u}_{(0,0)}$. These vector subspaces of $\tilde{u} \otimes \tilde{u}$ intersect in $\{0\}$, whence the condition is equivalent to $(1 \otimes E_i)R_0 = R_0(\tilde{K}_{-i} \otimes E_i)$ and $(E_i \otimes \tilde{K}_i)R_0 = R_0(E_i \otimes 1)$. Now

$$\begin{aligned} (1 \otimes E_i)R_0 &= \sum_{\mu, \rho} f(\mu, \rho) q^{-\langle \rho, i' \rangle} (K_{\mu+[i]} \otimes K_\rho) (\tilde{K}_{-i} \otimes E_i) \\ &\stackrel{!}{=} \sum_{\mu, \rho} f(\mu+[i], \rho) (K_{\mu+[i]} \otimes K_\rho) (\tilde{K}_{-i} \otimes E_i) = R_0(\tilde{K}_{-i} \otimes E_i) \end{aligned}$$

holds if and only if $f(\mu+[i], \rho) = f(\mu, \rho) q^{-\langle \rho, i' \rangle}$ for all μ, ρ . A similar argument shows that $(E_i \otimes \tilde{K}_i)R_0 = R_0(E_i \otimes 1)$ is equivalent to $f(\mu, \rho+[i]) = f(\mu, \rho) q^{-\langle \mu, i' \rangle}$ for all μ, ρ . The equations for F_i give the same conditions for f , with i replaced by $-i$, which are not really new.

(b) $(\Delta \otimes \text{id})R = R^{13}R^{23}$. We have

$$(\Delta \otimes \text{id})R = \sum_{\nu_1, \nu_2; \mu, \rho} f(\mu, \rho) (K_\mu \otimes K_{\mu-[\nu_1]} \otimes K_\rho) \bar{\Theta}_{\nu_1}^{13} \bar{\Theta}_{\nu_2}^{23}$$

and (summation over $\nu_1, \nu_2, \mu_1, \mu_2, \rho_1, \rho_2$)

$$\begin{aligned} R^{13}R^{23} &= \sum f(\mu_1, \rho_1) f(\mu_2, \rho_2) (K_{\mu_1} \otimes K_{\mu_2} \otimes K_{\rho_1+\rho_2}) q^{-\langle \rho_2, \nu'_1 \rangle} \bar{\Theta}_{\nu_1}^{13} \bar{\Theta}_{\nu_2}^{23} \\ &\stackrel{(26)}{=} \sum f(\mu_1, \rho_1) f(\mu_2 + [\nu_1], \rho_2) (K_{\mu_1} \otimes K_{\mu_2} \otimes K_{\rho_1+\rho_2}) \bar{\Theta}_{\nu_1}^{13} \bar{\Theta}_{\nu_2}^{23} \\ &= \sum f(\mu_1, \rho_1) f(\mu_2, \rho_2) (K_{\mu_1} \otimes K_{\mu_2-[\nu_1]} \otimes K_{\rho_1+\rho_2}) \bar{\Theta}_{\nu_1}^{13} \bar{\Theta}_{\nu_2}^{23}. \end{aligned}$$

Since the group-like elements are linearly independent, we have equality if and only if for all ρ, μ, μ' ,

$$\sum_{\rho_1 + \rho_2 = \rho} f(\mu, \rho_1) f(\mu', \rho_2) = \delta_{\mu, \mu'} f(\mu, \rho).$$

The other condition in (27) is equivalent to $(\text{id} \otimes \Delta)R = R^{13}R^{12}$.

(c) R is invertible. Assuming the conditions (25) for R , we have the following equivalent conditions:

- (i) R is invertible
- (ii) $(\varepsilon \otimes \text{id})R = 1$,
- (iii) $(\text{id} \otimes \varepsilon)R = 1$.

Proof. If R is invertible, R is a quasitriangular structure, and [5, Theorem VIII.2.4] implies (ii) and (iii), since the antipode of \tilde{u} is clearly invertible. It follows from (ii) (or (iii)) that $(S \otimes \text{id})R$ (or $(\text{id} \otimes S^{-1})R$) is an inverse of R (cf. proof of [5, Theorem VIII.2.4]). We have $(\varepsilon \otimes \text{id})\bar{\Theta} = (\text{id} \otimes \varepsilon)\bar{\Theta} = 1$ because the only non-zero contribution comes from $\bar{\Theta}_0 = 1 \otimes 1$. Direct calculation shows that $(\text{id} \otimes \varepsilon)R = (\varepsilon \otimes \text{id})R = 1$ is equivalent to (28). If there exists $l \in k \setminus \{0\}$ such that $f(\mu, 0) = f(0, \rho) = l$ for all μ, ρ then (27) implies for $\mu' = 0$,

$$\sum_{\rho_1 + \rho_2 = \rho} f(\mu, \rho_1) l = \delta_{\mu, 0} f(\mu, \rho) = \delta_{\mu, 0} l \Leftrightarrow \sum_{\rho_1} f(\mu, \rho_1) = \delta_{\mu, 0}.$$

■

Remark 8.12. Conditions (27) and (28) just mean that R_0 is a quasitriangular structure of the group algebra $k[\{K_\mu \mid \mu \in \tilde{Y}\}]$ by [8]. Then (26) is a compatibility condition.

Note. It is always possible to choose y_1, y_2 such that $y_2 = y_1$ and $|y_1|f: y_1 x y_1 \rightarrow k \setminus \{0\}$ is a bicharacter.

Another class of quasitriangular structures can be constructed from the above.

PROPOSITION 8.13. (a) *If a Hopf algebra H admits a quasitriangular structure $R \in H \otimes H$ then $\tau(R^{-1})$ is a quasitriangular structure, too, where $\tau: H \otimes H \rightarrow H \otimes H$, $x \otimes y \mapsto y \otimes x$ denotes the usual twist.*

(b) *Let $R_0 = \sum f(\mu, \rho) K_\mu \otimes K_\rho$ as in Theorem 8.11. Then $R_0^{-1} = \sum f(\mu, -\rho) K_\mu \otimes K_\rho = \sum f(-\mu, \rho) K_\mu \otimes K_\rho$. In particular, $f(\mu, \rho) = f(-\mu, -\rho)$ for all μ, ρ .*

(c) If f satisfies (26)–(28) then \tilde{u} admits the quasitriangular structures

$$R_0 \bar{\Theta} = \left(\sum_{\mu, \rho} f(\mu, \rho) (K_\mu \otimes K_\rho) \right) \left(\sum_{\nu} (-1)^{\text{tr } \nu} q_\nu^{-1} \sum \bar{b}^- \otimes (\bar{b}^*)^+ \right)$$

$$\tau(\Theta) \tau(R_0^{-1}) = \left(\sum_{\nu} (-1)^{\text{tr } \nu} q_\nu \sum b^+ \otimes b^{*-} \right) \left(\sum_{\mu, \rho} f(\mu, -\rho) K_\rho \otimes K_\mu \right).$$

Proof. (a) From [5, Theorem VIII.2.4(b)] we get $(S \otimes \text{id})R = R^{-1}$. Then direct calculation proves (25).

(b) We show that $\tilde{R}_0 = \sum f(\mu, -\rho) K_\mu \otimes K_\rho$ is an inverse of R_0 . We have

$$R_0 \tilde{R}_0 = \sum_{\substack{\mu_1 + \mu_2 = \mu \\ \rho_1 + \rho_2 = \rho}} f(\mu_1, \rho_1) f(\mu_2, -\rho_2) K_\mu \otimes K_\rho$$

and

$$\sum_{\substack{\mu_1 + \mu_2 = \mu \\ \rho_1 + \rho_2 = \rho}} f(\mu_1, \rho_1) f(\mu_2, -\rho_2) \stackrel{(27)}{=} \sum_{\rho_1 + \rho_2 = \rho} \delta_{\rho_1, -\rho_2} f(\mu, \rho_1) \stackrel{(28)}{=} \delta_{\rho, 0} \delta_{\mu, 0}.$$

This implies $R_0 \tilde{R}_0 = 1 \otimes 1$. Since R_0 and \tilde{R}_0 commute, \tilde{R}_0 is the inverse. A similar computation gives the second expression for the inverse. Combining both gives $f(\mu, \rho) = f(-\mu, -\rho)$.

(c) This follows from Remark 8.6. ■

PROPOSITION 8.14. Assume that there are no weak diagram antimorphisms of (I, \cdot) (e.g., if it is irreducible). Two quasitriangular structures $R_0 \bar{\Theta}$ and $R'_0 \bar{\Theta}$, where $R_0 = \sum f(\mu, \rho) K_\mu \otimes K_\rho$ and $R'_0 = \sum f'(\mu, \rho) K_\mu \otimes K_\rho$ are equivalent, i.e., there is a Hopf automorphism ϕ such that $(\phi \otimes \phi)(R_0 \bar{\Theta}) = R'_0 \bar{\Theta}$ if and only if there exist isomorphisms $\gamma_1: Y_1 \rightarrow Y'_1$, $\gamma_2: Y_2 \rightarrow Y'_2$ such that $f(\mu, \rho) = f'(\gamma_1(\mu), \gamma_2(\rho))$ and $\phi(K_\mu) = K_{\gamma_1(\mu)}$, $\phi(K_\rho) = K_{\gamma_2(\rho)}$ for all μ, ρ . The same holds for equivalence of $\tau(\Theta) \tau(R_0)$ and $\tau(\Theta) \tau(R'_0)$. Two quasitriangular structures of the types $R_0 \bar{\Theta}$ and $\tau(\Theta) \tau((R'_0)^{-1})$ are never equivalent.

Proof. Proposition 8.7 implies $(\phi \otimes \phi) \bar{\Theta} = \bar{\Theta}$ and $(\phi \otimes \phi) \tau(\Theta) = \tau(\Theta)$. Hence we need $(\phi \otimes \phi)(R_0) = R'_0$ in both cases which is equivalent to the conditions in the assertion. The last statement holds because $R_0 \bar{\Theta} \subset \tilde{u}^- \tilde{u}^0 \otimes \tilde{u}^+ \tilde{u}^0$, $\tau(\Theta) \tau((R'_0)^{-1}) \in \tilde{u}^+ \tilde{u}^0 \otimes \tilde{u}^- \tilde{u}^0$ and each automorphism maps \tilde{u}^\pm to \tilde{u}^\pm . ■

We give explicit examples.

THEOREM 8.15. Let $Y_1 = Y_2$ be subgroups of the group generated by I in \tilde{Y} , which contain the images of $[i] \in Y$ for all $i \in I$ and satisfy Condition

(29). Define a \mathbb{Z} -bilinear map $\mathbb{Z}[I] \times \mathbb{Z}[I] \rightarrow \mathbb{Q}$ by

$$i \diamond j := \frac{i \cdot j}{(1/2)(i \cdot i)(1/2)(j \cdot j)} = \frac{\langle i, j' \rangle}{(1/2)j \cdot j} = \frac{\langle j, i' \rangle}{(1/2)i \cdot i}.$$

Let g be the lowest common denominator of the numbers $\{\nu \diamond \tilde{\nu} \mid \nu, \tilde{\nu} \in Y_1\}$. Assume that there exists $\hat{q} \in k$ such that $\hat{q}^g = q$. Define $Z := \{\mu \in Y_1 \mid \forall \rho \in Y_1: \hat{q}^{g\mu \diamond \rho} = 1; K_\mu \text{ central}\}$, let $l = |Y_1/Z|$, and assume that $\text{char}(k)$ does not divide l . Then $\tilde{u} := \tilde{u}/(\{K_\mu - 1 \mid \mu \in Z\})$ has a quasitriangular structure as above where $f(\mu, \rho) = \hat{q}^{-g\mu \diamond \rho}/l$.

Proof. f is well-defined on $Y_1 \times Y_1$, because (29) holds. Since Z only contains elements μ such that $\hat{q}^{g\mu \diamond \rho} = 1$ for all $\rho \in Y_1$, and \diamond is symmetric, f is well-defined on $Y_1/Z \times Y_1/Z$. Since f is symmetric in its arguments, it suffices to prove the first conditions of (26), (27). For all $i \in I$ we have

$$f(\mu + [j], \rho) = \frac{\hat{q}^{-g\mu \diamond \rho} \hat{q}^{-g[j] \diamond \rho}}{l} = f(\mu, \rho) \hat{q}^{-g\langle \rho, j' \rangle} = f(\mu, \rho) q^{-\langle \rho, j' \rangle}.$$

Moreover

$$\begin{aligned} \sum_{\rho_1 + \rho_2 = \rho} f(\mu, \rho_1) f(\tilde{\mu}, \rho_2) &= \frac{1}{l^2} \sum_{\rho_1 + \rho_2 = \rho} \hat{q}^{-g\mu \diamond (\rho_1 + \rho_2) - g(\tilde{\mu} - \mu) \diamond \rho_2} \\ &= \frac{1}{l} f(\mu, \rho) \sum_{\rho_2} \hat{q}^{-g(\tilde{\mu} - \mu) \diamond \rho_2}. \end{aligned}$$

For all $\mu \in Y_1$, the map $Y_2 \rightarrow k \setminus \{0\}$, $\rho \mapsto \hat{q}^{-g\mu \diamond \rho}$ is a group character, hence (27) follows from $|Y_1| = |Y_2|$ under the condition that this character is non-trivial for $\mu \notin Z$. Let $\mu \in Y_1$. If K_μ is not central, then there exists $i \in I$ such that $E_i K_\mu \neq K_\mu E_i = q^{\langle \mu, i' \rangle} E_i K_\mu$ (or $F_i K_\mu \neq K_\mu F_i$), whence $\hat{q}^{g\mu \diamond [i]} \neq 1$. If K_μ is central and $\hat{q}^{g\mu \diamond \rho} = 1$ for all $\rho \in Y_1$, then $\mu \in Z$ and $K_\mu = 1$ in \tilde{u} . Condition (28) is automatically satisfied. ■

EXAMPLE 8.16. (a) The “standard” quasitriangular structure (e.g., [3, Proposition 9.3.9]). If $G \subset \tilde{Y}$ contains all central group-like elements of the form \tilde{K}_ν , Y_1 is generated by the images of $[i]$ in \tilde{Y}/G , and $\text{char}(k)$ does not divide $|Y_1|$, then $\tilde{u} = \tilde{u}/(\{K_\mu - 1 \mid \mu \in G\})$ has quasitriangular structures as above where

$$R_0 = \frac{1}{|Y_1|} \sum_{\nu, \tilde{\nu} \in Y_1} q^{-\nu \cdot \tilde{\nu}} \tilde{K}_\nu \otimes \tilde{K}_{\tilde{\nu}}.$$

(b) Assume that $\frac{i \cdot i}{2}$ is coprime to the order of q for all $i \in I$ and that G contains all central group-like elements of the form K_ν , $\nu \in \mathbb{Z}[I]$. Let Y_1 be the image of $\mathbb{Z}[I]$ in \tilde{Y}/G . Assume that $\text{char}(k)$ does not divide $|Y_1|$.

Then \tilde{u} has quasitriangular structures as above where

$$R_0 = \frac{1}{|Y_1|} \sum_{\nu, \tilde{\nu} \in Y_1} \hat{q}^{-g\nu \diamond \tilde{\nu}} K_\nu \otimes K_{\tilde{\nu}},$$

where g, \diamond are as in Theorem 8.15 and \hat{q} is a power of q .

Proof. (a) For all $i, j \in I$ we have $[i] \diamond [j] = \langle [i], j' \rangle = i \cdot j \in \mathbb{Z}$, hence $g = 1$. Moreover, $q^{2l_j \langle [i], j' \rangle} = (q_j^{2l_j})^{\langle j, i' \rangle} = 1$. We have $Z = \{0\}$ because \tilde{u} does not contain nontrivial central group-like elements of the form \tilde{K}_ν where $[\nu] \in Y_1$.

(b) Since $\frac{i \cdot i}{2}$ is coprime to the order of q , q_i has the same order as q and \hat{q} can be chosen as a power of q . Therefore $q^{2l_i} = 1$, and (29) is satisfied. \tilde{u} does not contain nontrivial central group-like elements of the form K_ν , $\nu \in \mathbb{Z}[I]$, whence $Z = \{0\}$. ■

Here we give a more explicit description of the quasitriangular structures from the classification in [4]. We recall the situation of the paper in our setting. The element “ \sqrt{q} ” in [4] is called q here. Its order is N as in [4]. We have the Cartan datum (I, \cdot) given by $I = \{i_1, \dots, i_{n-1}\}$, and

$$i_\alpha \cdot i_\beta = \begin{cases} 4 & \text{if } \alpha = \beta \\ -2 & \text{if } |\alpha - \beta| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover we have $Y = \mathbb{Z}[I]$, $K_i^N = 1$, and $q^8 \neq 1$. We will call a quasitriangular structure *minimal* in the sense of [11], that is, if there is no proper Hopf subalgebra A such that the quasitriangular structure is an element of $A \otimes A$. In particular, we have $Y_1 = \tilde{Y}/G$ for minimal quasitriangular structures. Denote the highest common divisor of two positive integers a, b by $\text{hcd}(a, b)$. The classification is given in [4, Theorem 4.1]:

THEOREM 8.17. *Let $Y_0 = 2\tilde{Y}$ be the subgroup generated by $[i]$ for $i \in I$. Let $\omega = i_1 + i_3 + \dots + i_{n-1}$ if n is even; let Y_ω be generated by Y_0 and ω and assume that $\text{char}(k)$ does not divide $|Y_\omega|$ and $|Y_0|$.*

(a) *If $n > 2$ and $\text{hcd}(n, N) = 1$, then \tilde{u} admits two non-isomorphic quasitriangular structures for $Y_1 = Y_2 = Y_0$ (being minimal if and only if N is odd).*

(b) *If $\text{hcd}(n, N) = 2$ and $\frac{N}{2}$ is odd then \tilde{u} admits two non-isomorphic not minimal quasitriangular structures for $Y_1 = Y_2 = Y_0$ and two non-isomorphic quasitriangular structures for $Y_1 = Y_2 = Y_\omega$ (being minimal if and only if $n = 2$).*

(c) If $\text{hcd}(n, N) = 2$ and $\frac{N}{2}$ is even then $\tilde{\mathfrak{u}}$ admits four non-isomorphic quasitriangular structures for $Y_1 = Y_2 = Y_\omega$ (being minimal if and only if $n = 2$).

Proof. We only show the existence. For each quasitriangular structure R , we consider the non-isomorphic quasitriangular structure $\tau(R^{-1})$, too (cf. Propositions 8.13 and 8.14).

(a) By [10, 4.3(c)] and $K_i^N = 1$ for all i , $\tilde{\mathfrak{u}}$ does not contain central group-like elements (except 1). If N is odd then it follows from $K_i = K_i^{(N+1)/2}$ that these quasitriangular structures are minimal.

(b) and (c) We show first that the quasitriangular structures for Y_ω are well-defined. We have

$$\omega \diamond \omega = \frac{n}{2} \in \mathbb{N}, \quad \omega \diamond [i_s] = \begin{cases} 2 & \text{if } s \text{ is odd,} \\ -2 & \text{if } s \text{ is even} \end{cases}$$

and $q^{2l[i] \diamond \omega} = 1$ since $q_i = q^2$. Hence the lowest common denominator g is equal to 1 and $f(\mu, \rho) = q^{-\mu \diamond \rho}$ for $\mu, \rho \in Y_1$. Now we show that $Z = \{0\}$, i.e., no central group-likes have to be divided out. Since we have $2\omega \in Y_0$ and Z is a group, it suffices to consider elements of the types $\omega + [\nu]$, where $\nu \in Y_0$ such that $\tilde{K}_{2\nu} = 1$ and $[\nu]$ where $\nu \in Y_0$. If $\frac{N}{2}$ is odd then \tilde{K}_ν has odd order, hence $\omega + [\nu] = \omega$, but

$$q^{\omega \diamond [i_1]} = q^2 \neq 1.$$

If $\frac{N}{2}$ is even then $\frac{n}{2}$ must be odd and

$$q^{(\omega + [\nu]) \diamond \omega N/2} = (-1)^{n/2} \neq 1$$

because $[\nu] \diamond \omega$ is even. Now we examine elements of the second type, in particular \tilde{K}_ν is central. We consider the system of linear equations in [10, 4.3(b)] for central group-like elements and write $N = 2^r t$ where $r \in \mathbb{N}_0$ and t is odd. By the Chinese remainder theorem it is enough to solve the system modulo 2^r and modulo t . But n is coprime to t , whence all entries in the solutions must be multiples of t and the order of a central group-like element is a power of 2. If there are no such elements except 1, we are done, e.g., if $\frac{N}{2}$ is odd, then all elements \tilde{K}_ν have odd order. Otherwise, $\frac{n}{2}$ is odd and we replace the element by a power of it, which has order 2. It is shown in the proof of Theorem 2.2 in [4] that it must be equal to $K_\omega^{N/2}$. But $q^{\omega N/2 \diamond \omega} = (-1)^{n/2} \neq 1$ since $\frac{n}{2}$ is odd.

Hence we have found all quasitriangular structures if $\frac{N}{2}$ is odd and it is clear that they are minimal if and only if $n = 2$. It remains to show that the quasitriangular structures are not equivalent if $\frac{N}{2}$ is even. By [10, 5.8] for each automorphism ϕ of $\tilde{\mathfrak{u}}$ there exists a diagram automorphism σ of I (either $\sigma = \text{id}$ or $\sigma(i_t) = i_{n-t}$ for all t) such that $\phi(K_i^2) = K_{\sigma(i)}^2$ for all $i \in I$. Moreover $\phi(K_i)K_{\sigma(i)}^{-1}$ is central group-like with square 1, whence $\phi(K_i) \in \{K_{\sigma(i)}, K_{\sigma(i)}K_\omega^{N/2}\}$ and $\phi(K_\omega) \in \{K_\omega, K_\omega^{N/2+1}\}$. In both cases, f remains unchanged because $\frac{N}{2}$ is even and $q^{(N/2+1)\omega \diamond (N/2+1)\omega} = q^{\omega \diamond \omega}$. ■

8.3. Ribbon Elements

We prove that all quasitriangular structures which have been considered in the previous subsection admit ribbon elements. We recall the definition of ribbon elements.

DEFINITION 8.18. Let H be a quasitriangular Hopf algebra with quasitriangular structure $R = \sum R^{(1)} \otimes R^{(2)}$. Define the element $u := \sum S(R^{(2)})R^{(1)}$. The algebra H is called a *ribbon Hopf algebra* if there exists an invertible central element $\mathbf{v} \in H$ (which will be called a *ribbon element*) such that

$$\mathbf{v}^2 = uS(u), \quad S(\mathbf{v}) = \mathbf{v}, \quad \varepsilon(\mathbf{v}) = 1, \quad \Delta(\mathbf{v}) = (\tau(R)R)^{-1}(\mathbf{v} \otimes \mathbf{v}).$$

We recall some properties of R and the element u of Definition 8.18.

LEMMA 8.19. Let $R = \sum R^{(1)} \otimes R^{(2)}$, u be as in Definition 8.18. Then

$$\begin{aligned} R^{-1} &= (S^{-1} \otimes \text{id})R = (\text{id} \otimes S)R \\ R &= (S \otimes S)R \end{aligned} \tag{30}$$

$$S^2(x) = uxu^{-1} \quad \text{for all } x \in H \tag{31}$$

$$\Delta(u) = (u \otimes u)(\tau(R)R) = (\tau(R)R)(u \otimes u), \quad \varepsilon(u) = 1$$

$$\Delta(S(u)) = (S(u) \otimes S(u))(\tau(R)R) = (\tau(R)R)(S(u) \otimes S(u))$$

$$u^{-1} = \sum S^{-1}(R^{(1)})S(R^{(2)}).$$

Proof. See [5, Propositions VIII.4.1 and VIII.4.5]. ■

LEMMA 8.20. Let R be a quasitriangular structure and $\check{R} = \tau(R^{-1})$. Compute the elements u and \check{u} from R and \check{R} as in Definition 8.18. Then

$$(a) \quad u = S(\check{u})^{-1}, \quad S(u) = \check{u}^{-1}.$$

(b) If \mathbf{v} is a ribbon element for R , then \mathbf{v}^{-1} is a ribbon element for \check{R} and vice versa.

Proof. (a) It suffices to prove $S(u) = \check{u}^{-1}$, because $S^2(u) = u$, $S^2(\check{u}) = \check{u}$, cf. (31). We have

$$\check{u}^{-1} = \sum S^{-2}(S(R^{(2)}))R^{(1)} = \sum S^{-1}(S(R^{(1)})R^{(2)}) = S^{-1}(u) = S(u).$$

(b) The conditions in Definition 8.18 can be checked using $\check{u}S(\check{u}) = S(u)^{-1}u^{-1} = (uS(u))^{-1}$ and $(\tau(\check{R})\check{R})^{-1} = \tau(R)R$. ■

We recall some facts about root systems using slightly modified notions of [6].

DEFINITION-LEMMA 8.21. For all $j \in I$ define the maps $(s_j)_Y: Y \rightarrow Y$ by $(s_j)_Y(\mu) = \mu - \langle \mu, j' \rangle j$ for $\mu \in Y$ and $(s_j)_X: X \rightarrow X$ by $(s_j)_X(\lambda) = \lambda - \langle j, \lambda \rangle j'$ for $\lambda \in X$. These maps induce actions of the Weyl group W on X and on Y : $(w, \lambda) \mapsto w_X(\lambda), (w, \mu) \mapsto w_Y(\mu)$. For all $\lambda \in X, \mu \in Y$ we have $\langle w_Y(\mu), \lambda \rangle = \langle \mu, w_X^{-1}(\lambda) \rangle$ [6, Paragraph 2.2.6]. Since (I, \cdot) is of finite type, $\mathbb{Z}[I]$ can be canonically imbedded into X and Y . Let W act on $\mathbb{Z}[I]$ via $w_X(i)' = w_X(i')$. Following [6, Paragraph 2.3.1], we define the sets $\tilde{\mathcal{R}}$ and \mathcal{R} of roots and coroots, respectively, as sets of all $\lambda \in X$ and $\mu \in Y$ where $\lambda = w_X(i'), \mu = w_Y(i)$ for some $i \in I, w \in W$. Now $\tilde{\mathcal{R}}$ and \mathcal{R} are finite subsets of $\sum_i \mathbb{Z}i' \subset X$ and $\sum_i \mathbb{Z}i \subset Y$. Let $\tilde{\mathcal{R}}^+ = \tilde{\mathcal{R}} \cap \mathbb{N}_0[I]$ and $\mathcal{R}^+ = \mathcal{R} \cap \mathbb{N}_0[I]$ be the sets of positive roots and coroots, respectively. Let $\gamma \in \mathbb{N}_0[I] \subset X$ be the sum of all positive roots. Then

(a) $\gamma \cdot i = i \cdot i$ for all $i \in I$.

(b) For each root $w_X(i)$ and the corresponding coroot $w_Y(i)$ we have $w_Y([i]) = [w_X(i)]$.

Proof. (a) This is a property of all root systems.

(b) Direct calculation shows that for all $\nu, \rho \in \mathbb{Z}[I]$ and $l \in I$ we have $(s_l)_X(\nu) \cdot \rho = \nu \cdot (s_l)_X^{-1}(\rho)$. Hence for each $j \in I$ we conclude

$$\begin{aligned} \langle w_Y([i]), j' \rangle &= \frac{i \cdot i}{2} \langle i, w_X^{-1}(j)' \rangle = i \cdot w_X^{-1}(j) \\ &= w_X(i) \cdot j = \langle [w_X(i)], j' \rangle. \end{aligned}$$

Since $w_Y([i])$ and $[w_X(i)]$ are in $\mathbb{Z}[I]$ and the Cartan matrix is invertible, they must be equal. ■

LEMMA 8.22. Let $\gamma \in \mathbb{Z}[I]$ be as in Definition-Lemma 8.21. Then for all $x \in \tilde{\mathfrak{t}}$

$$S^2(x) = \tilde{K}_\gamma^{-1} x \tilde{K}_\gamma.$$

Proof. It suffices to check it on the algebra generators, where it is an easy calculation. ■

THEOREM 8.23. All quasitriangular structures considered in the previous subsection admit ribbon elements.

Proof. By Lemma 8.20 it suffices to consider quasitriangular structures $R = R_0 \bar{\Theta}$, where $R_0 = \sum f(\mu, \rho) K_\mu \otimes K_\rho$. We write $\bar{\Theta} = \sum \bar{\Theta}^{(1)} \otimes \bar{\Theta}^{(2)}$. We use (30) and obtain

$$u = \sum f(\mu, \rho) K_\rho \bar{\Theta}^{(2)} S^{-1} (K_\mu \bar{\Theta}^{(1)}) = \sum f(\mu, \rho) K_\rho \bar{\Theta}^{(2)} S^{-1} (\bar{\Theta}^{(1)}) K_{-\mu}.$$

If a summand in $\bar{\Theta}^{(2)}$ is in \tilde{u}_ν^+ , then the corresponding summand in $\bar{\Theta}^{(1)}$ and of $S^{-1}(\bar{\Theta}^{(1)})$ belongs to $\tilde{u}_\nu^- \tilde{u}^0$. Hence every group-like element commutes with $\vartheta := \Sigma \bar{\Theta}^{(2)} S^{-1}(\bar{\Theta}^{(1)})$, and

$$u = \sum f(\mu, \rho) K_{\mu-\rho} \vartheta = \left(\sum f(\mu, \rho) K_{\mu+\rho} \right)^{-1} \vartheta$$

by Proposition 8.13(b); this proposition also implies

$$S\left(\sum f(\mu, \rho) K_{\mu+\rho}\right) = \sum f(\mu, \rho) K_{-\mu-\rho} = \sum f(\mu, \rho) K_{\mu+\rho}.$$

Hence

$$S(u) = \left(\sum S(f(\mu, \rho) K_{\mu+\rho}) \right)^{-1} S(\vartheta) = \left(\sum f(\mu, \rho) K_{\mu+\rho} \right)^{-1} S(\vartheta).$$

$u^{-1}S(u)$ is group-like by Lemma 8.19. Hence there exists $\mu \in Y$ such that $S(u) = K_\mu u$. We compute K_μ . There exists $\nu \in \mathbb{N}_0[I]$ with maximal trace $\text{tr } \nu$ such that $\tilde{\mathbf{f}}_\nu \neq \{0\}$, because $\tilde{\mathbf{f}}$ is finite-dimensional. Now $S^{-1}(\bar{\Theta}_\nu^{(1)}) = \check{\Theta}_\nu^{(1)} \tilde{K}_\nu$, where $\check{\Theta}_\nu^{(1)} \in \tilde{u}_\nu^-$, whence we have the summand $\tilde{K}_\nu \bar{\Theta}_\nu^{(2)} \check{\Theta}_\nu^{(1)}$ of ϑ . On the other hand, $S(\vartheta) = \Sigma \bar{\Theta}^{(1)} S(\bar{\Theta}^{(2)})$ and $S(\bar{\Theta}_\nu^{(2)}) = \check{\Theta}_\nu^{(2)} \tilde{K}_{-\nu}$, where $\check{\Theta}_\nu^{(2)} \in \tilde{u}_\nu^+$, whence we have the summand $\tilde{K}_{-\nu} \bar{\Theta}_\nu^{(1)} \check{\Theta}_\nu^{(2)}$ of $S(\vartheta)$. When we use the commutation formulae for elements of the $+$ and $-$ parts of \tilde{u} (cf. [7, 6.5(a2)] or [6, Corollary 3.1.9]), contributions of homogeneous components of lower trace of degree do not affect the contributions of degree ν . Hence $K_\mu = \tilde{K}_{-\nu}^2$. We choose γ as in Definition–Lemma 8.21. If we can show that $\tilde{K}_{-2\nu-2\gamma}$ is the square of a central group-like element K , then $\mathbf{v} := K \tilde{K}_\gamma u$ is a ribbon element—it is central because for all $x \in \tilde{u}$,

$$S^2(x) = uxu^{-1} = \tilde{K}_\gamma^{-1} x \tilde{K}_\gamma$$

by Lemmas 8.22 and 8.19, and

$$S(\mathbf{v}) = K^{-1} \tilde{K}_\gamma^{-1} S(u) = K^{-1} \tilde{K}_\gamma^{-1} \tilde{K}_{-\nu}^2 u = \mathbf{v};$$

the other conditions follow from Lemma 8.19. It remains to find K . It suffices to consider an irreducible Cartan datum, for if $I = I_1 \cup I_2$ where $i_1 \cdot i_2 = 0$ for all $i_1 \in I_1, i_2 \in I_2$, and if $\tilde{\mathbf{f}}_1, \tilde{\mathbf{f}}_2$ are the corresponding algebras and K_1 and K_2 central group-like elements (which are central in \tilde{u} , too) with the desired properties for $\tilde{\mathbf{f}}_1$ and $\tilde{\mathbf{f}}_2$, then it follows from $\tilde{\mathbf{f}} \cong \tilde{\mathbf{f}}_1 \otimes \tilde{\mathbf{f}}_2$ (cf. proof of [10, 2.14]) that $K = K_1 K_2$ is a central group-like element which satisfies the desired properties for $\tilde{\mathbf{f}}$.

(a) Assume that q fits to ${}_k \mathbf{f}$. We have a Poincaré–Birkhoff–Witt type basis of $\tilde{\mathbf{f}}$ (cf. [10, Subsections 1.4 and 2.9]) consisting of elements b_c ,

where $c_{i_l} < l_{i_l}$ for all l , where l_{i_l} is the order of $q_{i_l}^2$. The degree of

$$T'_{i_1, e} T'_{i_2, e} \cdots T'_{i_{l-1}, e} (E_{i_l}^{(c_l)})$$

for $c_l = 1$ is $\alpha_l := (s_{i_1})_X \cdots (s_{i_{l-1}})_X(i_l)$, where α_l runs through all positive roots in $\tilde{\mathcal{R}}^+$. Now $c_l < l_{i_l} =: l_{\alpha_l}$ (this is independent of the actual choice of reduced expression of the positive coroot α_l). Hence the maximal degree is

$$\nu = \sum_{\alpha \in \tilde{\mathcal{R}}^+} \alpha(l_\alpha - 1).$$

Therefore $2\gamma + 2\nu = \sum_{\alpha \in \tilde{\mathcal{R}}^+} 2l_\alpha \alpha$, and $\tilde{K}_{-2\nu-2\gamma} = 1$ because for each positive root $\alpha = w_X(i)$ we have $\tilde{K}_\alpha^{2l_\alpha} = K_{w_Y(i)}^{i \cdot il_i}$ by Definition–Lemma 8.21(b) and because for $w_Y(i) = (s_{i_1})_Y \cdots (s_{i_{l-1}})_Y(i_l)$ we have

$$K_{w_Y(i)}^{i \cdot il_i} = T'_{i_1, e} \cdots T'_{i_{l-1}, e} (K_i^{i \cdot il_i}) = 1.$$

Therefore $K = 1$ is an appropriate choice.

(b) If q does not fit to ${}_k \mathbf{f}$, then [10, 2.13] applies. Here $\nu = 2i + 2j$ and $\tilde{K}_{-\nu-\gamma}$ is central because

$$q^{-(\nu+\gamma) \cdot i} = q^{-2i \cdot i} = 1, \quad q^{-(\nu+\gamma) \cdot j} = q^{-2j \cdot j} = 1.$$

Therefore K can be chosen as $\tilde{K}_{-\nu-\gamma}$. ■

COROLLARY 8.24. *The Frobenius–Lusztig kernel for sl_n considered in [4] and Theorem 8.17 and the quotients considered in Example 8.16 are ribbon Hopf algebras for all quasitriangular structures of the type described in Theorem 8.11 and Proposition 8.13.*

8.4. Quasitriangular Structures for ${}_k \mathbf{U}$

We show that ${}_k \mathbf{U}$ becomes a topological quasitriangular Hopf algebra (cf. [3, remarks after 9.3.9]). We complete the tensor products ${}_{\mathcal{A}} \mathbf{U} \otimes {}_{\mathcal{A}} \mathbf{U}$ and ${}_k \mathbf{U} \otimes {}_k \mathbf{U}$ in an analogous way as described in [6, 4.1.1] and denote them by $({}_{\mathcal{A}} \mathbf{U} \otimes {}_{\mathcal{A}} \mathbf{U})^\wedge$ and $({}_k \mathbf{U} \otimes {}_k \mathbf{U})^\wedge$, respectively. In [10, 2.9] elements $b_{\mathbf{c}}, \tilde{b}_{\mathbf{c}} \in \mathbf{f}$ for $\mathbf{c} \in \mathbb{N}_0^n$ (where n depends on the Cartan datum) have been considered where $(b_{\mathbf{c}}, \tilde{b}_{\tilde{\mathbf{c}}}) = \delta_{\mathbf{c}, \tilde{\mathbf{c}}}$ for all $\mathbf{c}, \tilde{\mathbf{c}} \in \mathbb{N}_0^n$. Therefore for $\nu \in \mathbb{N}_0[I]$ the sums

$$\Theta_\nu = (-1)^{\text{tr } \nu} v_\nu \sum_{\mathbf{c}: b_{\mathbf{c}} \in \mathbf{f}_\nu} b_{\mathbf{c}}^- \otimes \tilde{b}_{\mathbf{c}}^+$$

(recall that $v_\nu = \prod_i v_i^{\nu_i} = v^{\text{tr } [\nu]}$ for $\nu = \sum_i \nu_i i \in \mathbb{Z}[I]$) are summands of a Quasi- \mathcal{R} -matrix $\Theta = \sum_\nu \Theta_\nu$ of the completion $({}_{\mathcal{A}} \mathbf{U} \otimes {}_{\mathcal{A}} \mathbf{U})^\wedge$ as in [6, 4.1.2]. In particular, the properties shown in [6, 4.1.2–3] and analogues of Proposi-

tion 8.5 are valid. After tensoring with k over \mathcal{A} , we get a Quasi- \mathcal{A} -Matrix of ${}_k\mathbf{U}$ which will be denoted by Θ again and satisfies analogous properties. The following theorem can be proved in a similar way as Theorem 8.11 (note that ${}_k\mathbf{U}$ is generated as algebra by $E_i^{(t)}, F_i^{(t)}, K_\mu$ for $i \in I, t \in \mathbb{N}_0, \mu \in Y$).

THEOREM 8.25. *Let Y_1, Y_2 be as in Theorem 8.11. Let $R_0 = \sum_{\mu, \rho} f(\mu, \rho) K_\mu \otimes K_\rho$ where $f: Y_1 \otimes Y_2 \rightarrow k$ is any map. The product $R_0 \Theta$ is a quasitriangular structure of ${}_k\mathbf{U}$ if and only if f satisfies the conditions (26)–(28).*

Note added in proof. If the Carban datum is not of finite type than $\hat{\mathbf{f}}$ and $\hat{\mathbf{u}}$ are always infinite dimensional.

ACKNOWLEDGMENTS

The author thanks his supervisor Professor H.-J. Schneider for his excellent guidance and for fruitful discussions and Professor N. Andruskiewitsch and Professor M. Takeuchi for many hints. He thanks the referee for his very useful and valuable comments, suggestions, and corrections, in particular in Proposition 6.1.

REFERENCES

1. N. Andruskiewitsch, Notes on extensions of Hopf algebras, *Canad. J. Math.* **48** (1996), 3–42 [appendix by N. Andruskiewitsch and H.-J. Schneider].
2. W. Chin and I. M. Musson, The coradical filtration for quantized enveloping algebras, *J. London Math. Soc.* **53** (1996), 50–62.
3. V. Chari and A. Pressley, “A Guide to Quantum Groups,” Cambridge Univ. Press, Cambridge, UK, 1994.
4. S. Gelaki and S. Westreich, On the quasitriangularity of $U_q(sl_n)'$, preprint, 1995.
5. C. Kassel, “Quantum Groups,” Graduate Texts in Math, Vol. 155, Springer-Verlag, New York, 1995.
6. G. Lusztig, “Introduction to Quantum Groups,” Prog. Math., Vol. 110, Birkhäuser, Boston, 1993.
7. G. Lusztig, Quantum groups at roots of 1, *Geom. Dedicata* **35** (1990), 89–114.
8. S. Majid, “Foundations of Quantum Group Theory,” Cambridge Univ. Press, Cambridge, UK, 1995.
9. S. Montgomery, “Hopf Algebras and Their Action on Rings,” CBMS Lecture Notes, Vol. 82, Amer. Math. Soc., Providence, RI, 1993.
10. E. Müller, Some topics on Frobenius–Lusztig kernels, I, *J. Algebra* **206** (1998), 624–658.
11. D. E. Radford, Minimal quasitriangular Hopf algebras, *J. Algebra* **157** (1993), 281–315.
12. E. Taft and R. Wilson, On antipodes in pointed Hopf algebras, *J. Algebra* **29** (1974), 27–32.
13. M. Takeuchi, The coradical filtration for the quantum algebra U , preprint, April 1994.
14. M. Takeuchi, Some topics on $GL_q(n)$, *J. Algebra* **147** (1992), 379–410.
15. M. Takeuchi, q -Representations of quantum groups, in “Conf. Proc.,” Vol. 16, Canad. Math. Soc., 1995.